

Selberg's Method and the Multiplicities of the Zeroes of the Riemann Zeta-Function

by

Timothy S. TRUDGIAN

(Received February 22, 2011)

(Revised May 17, 2011)

Dedicated to Professor Akio Fujii who had the patience to listen to my presentation when he might have otherwise had lunch.

Abstract. This article shows how the approximation of $S(T)$ by a Dirichlet polynomial, work begun by Selberg, can be used to recover bounds on the multiplicities of the zeroes of $\zeta(s)$.

1. Introduction

This paper concerns the properties of the moments of the function $S(T)$, the argument of the Riemann zeta-function; for some basic properties the reader is invited to examine [9, Ch. 9]. The function $S(T)$ is intimately connected with $N(T)$, the number of non-trivial zeroes¹ of $\zeta(\sigma + it)$ for $0 < t < T$ through the following formula

$$N(T) = \frac{T}{2\pi} \log \frac{T}{2\pi} - \frac{T}{2\pi} + \frac{7}{8} + S(T) + O(T^{-1}), \quad (1)$$

where the $O(T^{-1})$ term is continuous. This continuity of the error term is essential, since it enables one to infer, by (1), that when T passes over a zero of multiplicity k , which is counted k times in $N(T)$, the function $S(T)$ increases by k . Results on the rate of growth of $S(T)$ therefore provide information as to the likelihood of zeroes of large multiplicity. Let $M(T)$ denote the maximum multiplicity of a zero between T and $2T$. Then

$$S(T) = O(\log T) \implies M(T) = O(\log T), \quad (2)$$

and on the Riemann hypothesis

$$S(T) = O\left(\frac{\log T}{\log \log T}\right) \implies M(T) = O\left(\frac{\log T}{\log \log T}\right). \quad (3)$$

Equation (2) is due to von Mangoldt and (3) is due to Littlewood (see, e.g. [9, p. 214 and p. 350]). This reliance of results on $S(T)$ to produce bounds for $M(T)$ should not be pushed too far. Selberg [8] showed that

$$S(T) = \Omega_{\pm}\left(\frac{(\log T)^{1/3}}{(\log \log T)^{7/3}}\right),$$

¹Here, and hereafter, *zeroes* always refers to non-trivial zeroes of the zeta-function.

whereas it is believed that all the zeroes of the zeta-function should be simple. Nevertheless, one can pursue the connexion between moments of $S(T)$ and zeroes of high multiplicity. Throughout this paper, A denotes a positive constant, not necessarily the same at each occurrence.

2. Selberg's Approximation

Selberg proved the following theorem in [8, p. 39, Thm 4].

THEOREM 2.1 (Selberg). *If $T^\alpha < H \leq T$, where α is fixed and $\frac{1}{2} < \alpha \leq 1$, and, for m a positive integer, $T^{\frac{\alpha-1}{m}} \leq x \leq H^{\frac{1}{m}}$, then*

$$\int_T^{T+H} \left| S(t) + \frac{1}{\pi} \sum_{p \leq x} \frac{\sin(t \log p)}{\sqrt{p}} \right|^{2m} dt \leq c(m)H,$$

where $c(m)$ depends on m but not on T .

The factor $c(m)$ has been improved over the years: in Selberg's paper [*op. cit.*] it was not calculated explicitly. Fujii [3] calculated that $c(m) \leq (Am)^{4m}$, and this result, which was also proved by Ghosh [4], follows more or less directly from Selberg's original arguments. Tsang [11] showed that $c(m) \leq (Am)^{2m}$, where the improvement comes from a repeated application of Selberg's density theorem, specifically Lemma 5.2 of [10]. Karatsuba and Korolev [6] placed a bound on the explicit constant in the result of Tsang, to show that $c(m) < (\epsilon^{-3} e^{37} \pi^{-2} m^2)^m$, for any fixed positive $\epsilon < 0.001$. A different approach is due to Goldston [5]. On the assumption of the Riemann hypothesis and introducing a different weight to that used by Selberg, Goldston showed that $c(m) \geq (A \log m)^m$.

3. The connexion with multiplicities

Define $N_j(T)$ to be the number of zeroes of multiplicity j at height T . Fujii [2] showed that

$$\frac{N_j(2T) - N_j(T)}{N(2T) - N(T)} \ll e^{-A\sqrt{j}}, \quad (4)$$

by using $c(m) \leq (Am)^{4m}$ in Theorem 2.1. Later Fujii [1] proved (4) with a bound of e^{-Aj} ; an explicit version of this improvement was given by Korolev [7]. In general, if one can show that $c(m) \leq (Am)^{\alpha m}$ for some α , then one can show² that

$$\frac{N_j(2T) - N_j(T)}{N(2T) - N(T)} \ll \exp(-Aj^{2/\alpha}). \quad (5)$$

Thus if there is one zero of multiplicity j between T and $2T$ one can use (5) to show that

$$(T \log T)^{-1} \ll \exp(-Aj^{2/\alpha}),$$

²This is implicit in the work of Fujii [3]—I thank the referee for making this known to me.

so that

$$j \ll (\log T)^{\alpha/2}.$$

Thus Tsang's result (viz. $\alpha = 2$) recovers the best unconditional bound for $M(T)$. This shows that any improvement on the constant $c(m)$ is likely to be difficult since $c(m) \leq (Am)^{\alpha m}$ for $\alpha < 2$ would surpass that which is currently provable on the assumption of the Riemann hypothesis—cf. (3).

3.1. Concluding question

The problem: assume the Riemann hypothesis and improve the bound on $c(m)$ to show that

$$M(T) \ll \frac{\log T}{\log \log T}.$$

This, being (3), is already known, but it would be of interest to see how much could be recovered unconditionally using this new approach.

Acknowledgment. I wish to thank the referee for his or her helpful suggestions.

References

- [1] A. Fujii, *On the uniformity of the distribution of the zeros of the Riemann zeta function*, Journal für die reine und angewandte Mathematik, **302**, 1978, 167–205.
- [2] A. Fujii, *On the distribution of the zeros of the Riemann Zeta function in short intervals*, Bulletin of the American Mathematical Society, **81**, 1975, 139–142.
- [3] A. Fujii, *On the Zeros of Dirichlet L-functions, I*, Transactions of the American Mathematical Society, **196**, 1974, 225–235.
- [4] A. Ghosh, *On Riemann's zeta function—Sign changes of $S(t)$ in “Recent Progress in Analytic Number Theory”*, Academic Press, London, 1981, 25–46.
- [5] D. A. Goldston, *On the Function $S(T)$ in the theory of the Riemann Zeta-Function*, Journal of Number Theory, **27**, 1987, 149–177.
- [6] A. A. Karatsuba and M. A. Korolev, *Behaviour of the argument of the Riemann zeta-function on the critical line*, Russian Mathematical Surveys, **61**, (3), 2006, 389–482.
- [7] M. A. Korolev, *On multiple zeros of the Riemann zeta function*, Izvestiya: Mathematics, **70**, (3), 2006, 427–446.
- [8] A. Selberg, *Contributions to the theory of the Riemann zeta-function*, Arch. for Math. og Naturv. B, **48**, (5), 1946, 89–155.
- [9] E. C. Titchmarsh, *The Theory of the Riemann zeta-function* Oxford University Press, Oxford, 2nd edition, 1986.
- [10] K.-M. Tsang, *The Distribution of the Values of the Riemann Zeta-Function*, PhD thesis, Princeton University, 1984.
- [11] K.-M. Tsang, *Some Omega-theorems for the Riemann zeta-function*, Acta Arithmetica, 1986, XLVI, 369–395.

Timothy S. Trudgian
University of Lethbridge
Canada
e-mail: tim.trudgian@uleth.ca